

Anti-PPAR gamma 兔多克隆抗体

K133492 -20°C


产品信息

| | |
|---------------|--|
| 蛋白质全称 | 过氧化物酶体增殖物激活受体γ |
| 别名 | PPARG, CIMT1, GLM1, NR1C3, PPARG1, PPARG2, PPARGgamma, peroxisome proliferator activated receptor gamma, PPAR-gamma, Nuclear receptor subfamily 1 group C member 3, Pparg, Nr1c3 |
| 免疫原 | 小鼠 PPAR gamma 重组蛋白 |
| Uniprot ID | P37238 |
| 抗体亚型 | IgG |
| 纯化方式 | 亲和纯化 |
| 预测分子量 / 观测分子量 | 58 kDa / 50-60 kDa |

应用

WB 免疫印迹

物种: 人, 小鼠, 大鼠

稀释比: 1: 800-1: 1500

阳性样品: 肝

背景

This gene encodes a member of the peroxisome proliferator-activated receptor (PPAR) subfamily of nuclear receptors. PPARs form heterodimers with retinoid X receptors (RXRs) and these heterodimers regulate transcription of various genes. Three subtypes of PPARs are known: PPAR-alpha, PPAR-delta, and PPAR-gamma. The protein encoded by this gene is PPAR-gamma and is a regulator of adipocyte differentiation. Additionally, PPAR-gamma has been implicated in the pathology of numerous diseases including obesity, diabetes, atherosclerosis and cancer.

储存条件

储存条件: 自收到之日起 -20°C下储存一年, 避免反复冻融。
储存缓冲液: 含 0.15% ProClin300 防腐剂, 100 μg/mL 牛血清白蛋白和 50%甘油的磷酸盐缓冲液。

图像



WB 检测 PPAR gamma 蛋白(货号 K133492).
 样品: 经 RIPA 裂解液(货号 KSG2002)处理的蛋白质.
 封闭: 3%脱脂牛奶(货号 KSGC310001)溶于 TBST 溶液,
 室温孵育 1 小时.
 一抗: 1: 1000 稀释, 4°C 孵育过夜.
 二抗: HRP 标记山羊抗兔 IgG (H+L) (货号 KB69909), 1:
 5000 稀释, 室温孵育 1 小时.

For Research Use Only!
Ver. No.: V1-202603