

# Anti-Factor VII 兔多克隆抗体

K1339888 -20°C



## 产品信息

蛋白质全称	凝血因子 VII
别名	Eptacog alfa, F7, Proconvertin, SPCA, Factor VII, Serum prothrombin conversion accelerator
免疫原	小鼠 Factor VII 重组蛋白
Uniprot ID	P70375
抗体亚型	IgG
纯化方式	亲和纯化
预测分子量 / 观测分子量	50 kDa / 50 kDa

## 应用

### WB 免疫印迹

物种: 小鼠, 大鼠

稀释比: 1: 300-1: 600

阳性样品: 肝


## 背景

F7(Coagulation factor VII) is also named as serum prothrombin conversion accelerator(SPCA). It is the key triggering enzyme of the extrinsic coagulation pathway. Hepatocytes are the predominant source of F7 production under normal physiologic conditions, although F7 is also expressed by monocytes/macrophages in inflammation and atherosclerosis. This protein has 2 isoforms produced by alternative splicing.

## 储存条件

**储存条件:** 自收到之日起 -20°C下储存一年, 避免反复冻融。  
**储存缓冲液:** 含 0.15% ProClin300 防腐剂, 100 µg/mL 牛血清白蛋白和 50%甘油的磷酸盐缓冲液。

## 图像



WB 检测 Factor VII 蛋白(货号 K1339888).  
 样品: 经 RIPA 裂解液(货号 KSG2002)处理的蛋白质.  
 封闭: 3%脱脂牛奶(货号 KSGC310001)溶于 TBST 溶液,  
 室温孵育 1 小时.  
 一抗: 1: 500 稀释, 4°C 孵育过夜.  
 二抗: HRP 标记山羊抗兔 IgG (H+L) (货号 KB69909), 1:  
 5000 稀释, 室温孵育 1 小时.

For Research Use Only!

Ver. No.: V1-202603