

Anti-MCM-BP 兔多克隆抗体

K1342540 -20°C



产品信息

蛋白质全称	微型染色体维持复合体结合蛋白
别名	C10orf119, MCM-BP, MCMBP
免疫原	小鼠 MCM-BP 重组蛋白
Uniprot ID	Q8R3C0
抗体亚型	IgG
纯化方式	亲和纯化
预测分子量 / 观测分子量	73 kDa / 73 kDa

应用

WB 免疫印迹

物种：人
 稀释比：1: 300-1: 600
 阳性样品：肝

背景

Associated component of the MCM complex that acts as a regulator of DNA replication. Binds to the MCM complex during late S phase and promotes the disassembly of the MCM complex from chromatin, thereby acting as a key regulator of pre-replication complex (pre-RC) unloading from replicated DNA. Can dissociate the MCM complex without addition of ATP; probably acts by destabilizing interactions of each individual subunits of the MCM complex. Required for sister chromatid cohesion (By similarity).

储存条件

储存条件：自收到之日起 -20°C下储存一年，避免反复冻融。
储存缓冲液：含 0.15% ProClin300 防腐剂, 100 μg/mL 牛血清白蛋白和 50%甘油的磷酸盐缓冲液。

图像



WB 检测 MCM-BP 蛋白(货号 K1342540).
 样品: 经 RIPA 裂解液(货号 KSG2002)处理的蛋白质.
 封闭: 3%脱脂牛奶(货号 KSGC310001)溶于 TBST 溶液,
 室温孵育 1 小时.
 一抗: 1: 300 稀释, 4°C 孵育过夜.
 二抗: HRP 标记山羊抗兔 IgG (H+L) (货号 KB69909), 1:
 5000 稀释, 室温孵育 1 小时.

For Research Use Only!

Ver. No.: V1-202603