

# Anti-Syk 兔多克隆抗体

K135367 -20°C



## 产品信息

蛋白质全称	酪氨酸蛋白激酶 SYK
别名	Syk, SYK, p72-Syk, spleen tyrosine kinase, spleen associated tyrosine kinase
免疫原	小鼠 Syk 重组蛋白
Uniprot ID	Q99MN1
抗体亚型	IgG
纯化方式	亲和纯化
预测分子量 / 观测分子量	72 kDa / 75 kDa

## 应用

### WB 免疫印迹

物种: 人, 小鼠, 大鼠

稀释比: 1: 500-1: 1000

阳性样品: 脾


## 背景

Spleen tyrosine kinase, also known as Syk, is an enzyme encoded by the SYK gene. SYK, along with Zap-70, is a member of the Syk family of tyrosine kinases. These non-receptor cytoplasmic tyrosine kinases share a characteristic dual SH2 domain separated by a linker domain. However, activation of SYK relies less on phosphorylation by Src family kinases than Zap-70. While Syk and Zap-70 are primarily expressed in hematopoietic tissues, there is expression of Syk in a variety of tissues.

## 储存条件

**储存条件:** 自收到之日起 -20°C下储存一年, 避免反复冻融。  
**储存缓冲液:** 含 0.15% ProClin300 防腐剂, 100 µg/mL 牛血清白蛋白和 50%甘油的磷酸盐缓冲液。

## 图像



WB 检测 Syk 蛋白(货号 K135367).

样品: 经 RIPA 裂解液(货号 KSG2002)处理的蛋白质。  
 封闭: 3%脱脂牛奶(货号 KSGC310001)溶于 TBST 溶液, 室温孵育 1 小时。  
 一抗: 1: 500 稀释, 4°C 孵育过夜。  
 二抗: HRP 标记山羊抗兔 IgG (H+L) (货号 KB69909), 1: 5000 稀释, 室温孵育 1 小时。

For Research Use Only!

Ver. No.: V1-202603