

# Anti-VASH2 兔多克隆抗体

K1336006 -20°C



## 产品信息

蛋白质全称	微管蛋白酪氨酸羧基酶 2
别名	Vasohibin-2, Vasohibin-like protein, Vash2, Vashl
免疫原	小鼠 VASH2 重组蛋白
Uniprot ID	Q8C5G2
抗体亚型	IgG
纯化方式	亲和纯化
预测分子量 / 观测分子量	40 kDa / 43 kDa

## 应用

### WB 免疫印迹

物种：小鼠，大鼠

稀释比：1: 500-1: 1000

阳性样品：肝

## 背景

VASH2, also named as vasohibin 2, is an important pro-angiogenesis factor in solid tumor. It has been reported that VASH2 is expressed in mononuclear cells mobilized from bone marrow to promote angiogenesis. VASH2 also plays a key role in axon formation. VASH2 is localized as a nuclear type (with 311 amino acid residues) and cytoplasmic type (with 355 amino acid residues and low abundance). Cytoplasmic VASH2 is associated with carcinoma angiogenesis, while nuclear VASH2 may be associated with cell proliferation.

## 储存条件

**储存条件：**自收到之日起 -20°C 下储存一年，避免反复冻融。  
**储存缓冲液：**含 0.15% ProClin300 防腐剂, 100 μg/mL 牛血清白蛋白和 50% 甘油的磷酸盐缓冲液。

## 图像



WB 检测 VASH2 蛋白(货号 K1336006).  
 样品: 经 RIPA 裂解液(货号 KSG2002)处理的蛋白质.  
 封闭: 3%脱脂牛奶(货号 KSGC310001)溶于 TBST 溶液,  
 室温孵育 1 小时.  
 一抗: 1: 500 稀释, 4°C 孵育过夜.  
 二抗: HRP 标记山羊抗兔 IgG (H+L) (货号 KB69909), 1:  
 5000 稀释, 室温孵育 1 小时.

For Research Use Only!

Ver. No.: V1-202603