

Anti-Cip4 兔多克隆抗体

K1340872 -20°C



产品信息

蛋白质全称	Cdc42 互作蛋白 4
别名	CIP4, HSTP, Protein Felic, Salt tolerant protein, STOT, STP, TR interacting protein 10, TRIP-10, TRIP10
免疫原	小鼠 Cip4 重组蛋白
Uniprot ID	Q8CJ53
抗体亚型	IgG
纯化方式	亲和纯化
预测分子量 / 观测分子量	68 kDa / 80 kDa

应用

WB 免疫印迹

物种: 人, 小鼠

稀释比: 1: 1000-1: 2000

阳性样品: Hep G2 细胞, 肾


背景

TRIP10 (thyroid hormone receptor interactor 10), also known as CIP4 (Cdc42-interacting protein 4), is a novel adaptor protein which may be involved in plasma membrane remodeling. CIP4 has been reported to negatively regulate glucose uptake by promoting GLUT4 endocytosis in muscle cells. At least five splice variants for human CIP4 have been identified, including CIP4a (CIP4), CIP4b (FELIC), CIP4c, CIP4d (CIP4/2). Western blot analysis usually demonstrates a major band around 75 kDa and a weaker band about 85 kDa.

储存条件

储存条件: 自收到之日起 -20°C下储存一年, 避免反复冻融。
储存缓冲液: 含 0.15% ProClin300 防腐剂, 100 µg/mL 牛血清白蛋白和 50%甘油的磷酸盐缓冲液。

图像



WB 检测 Cip4 蛋白(货号 K1340872).
 样品: 经 RIPA 裂解液(货号 KSG2002)处理的蛋白质.
 封闭: 3%脱脂牛奶(货号 KSGC310001)溶于 TBST 溶液, 室温孵育 1 小时.
 一抗: 1: 1000 稀释, 4°C 孵育过夜.
 二抗: HRP 标记山羊抗兔 IgG (H+L) (货号 KB69909), 1: 5000 稀释, 室温孵育 1 小时.

For Research Use Only!

Ver. No.: V1-202603